arXiv:hep-ph/0205306v2 28 May 2002

T violation in $B \to K^* \ell^+ \ell^-$ from SUSY

Chuan-Hung Chen[a,b] and C. Q. Geng[c,d]
[a]Department of Physics, National Cheng Kung University Tainan, Taiwan, Republic of China
[b]Institute of Physics, Academia Sinica, Taipei, Taiwan 115, Republic of China
[c]Department of Physics, National Tsing Hua University, Hsinchu, Taiwan, Republic of China
[d]Theory Group, TRIUMF, 4004 Wesbrrok Mall, Vancouver, B.C. V6T 2A3, Canada

We propose to use T-odd momentum correlations as physical observables to study T violation in $B \to K^* l^+ l^-$ $(l = e, \mu)$ decays. We show that these observables are zero in the standard model but sizable at the level of 10% in the supersymmetric model. These large T violating effects are measurable at the B-factories in KEK and SLAC and future hadron colliders. We also point out that the T violating effects are insensitive to nonperturbative QCD contributions.

PACS 13.20.He, 11.30.Er, 12.60.Jv, 12.38.Bx

The study of flavor change neutral current (FCNC) in B decays has an enormous progress since the CLEO observation [1] of the radiative $b \to s\gamma$ decay. Recently, the process of $B \to K\mu^+\mu^-$ has been observed [2] at the Belle detector in the KEKB e^+e^- storage ring with the branching ratio of $(7.5^{+2.5}_{-2.1} \pm 0.9) \times 10^{-7}$, while the SM expectation is around 5×10^{-7} [3]. Experimental searches at the B-factories for $B \to K^* l^+ l^-$ are also within the theoretical predicted ranges [5].

Via B decays such as $B \to J/\Psi\, K$, we can test the origin of CP violation (CPV) in the SM which is the consequence of the CKM quark mixing matrix [4]. CP asymmetries (CPAs) in B decays are usually defined by $a_{CP} \propto \Gamma - \bar{\Gamma}$ and $A_{CP}(t) \propto \Gamma(t) - \bar{\Gamma}(t)$. The former case, called direct CPA or CP-odd observable, needs both weak CP violating and strong phases, while the latter one of the time dependent CPA contains not only a non-zero CP-odd phase but also the $B - \bar{B}$ mixing. We note that the present world average for $a_{CP}^{\Psi K}$ is 0.79 ± 0.12 [6,7] comparing with the SM prediction of 0.70 ± 0.10 [7].

To study CPV, one can also define some other useful observables by the momentum correlations. For example, in a three-body decay, the simplest ones are the triple correlations of $\vec{s} \cdot (\vec{p}_i \times \vec{p}_j)$ [8], where $\vec{s}$ is the spin carried by one of outgoing particles and $\vec{p}_i$ and $\vec{p}_j$ denote any two independent momentum vectors. Clearly, these triple momentum correlations are T-odd observables since they change sign under the time reversal (T) transformation of $t \to -t$. In terms of the CPT invariant theorem, T violation (TV) implies CPV. Therefore, studying of T-odd observables can also help us to understand the origin of CPV. We note that the T violating observables such as the above triple correlations do not require strong phases. In the decays of $B \to K^* l^+ l^-$ $(l = e, \mu$, and $\tau)$, the spin s can be the polarized lepton, s_l, or the K^* meson, $\epsilon^*(\lambda)$. Considering the polarized lepton, as known, the polarization is always associated with the lepton mass, and thus we expect that the T violating effects are suppressed and less than 1% for the light lepton modes [9]. Although the τ mode can escape from the suppression, the corresponding branching ratio (BR) which is $O(10^{-7})$ is about one order smaller than those of e and μ modes.

It is known that CPAs such as a_{CP} in $B \to K^* l^+ l^-$ are small even with weak phases of $O(1)$ due to the smallness of strong phases [10]. In this paper, we concentrate on the possibility of having large T-odd terms such as $\vec{\epsilon}_{K^*}(\lambda) \cdot (\vec{p}_{l^+} \times \vec{p}_{K^*}) \propto \varepsilon_{\mu\nu\alpha\beta} q^\mu \epsilon^{*\nu}(\lambda) p_l^\alpha P^\beta$ in the decays of $B \to K^* l^+ l^-$ $(l = e, \mu)$, and for simplicity we set $m_l = 0$. We will show that these types of T violation are zero in the SM but they could be sizable in new physics such as the theories with SUSY.

The effective Hamiltonian of $b \to s l^+ l^-$ is given by [11]

$$\mathcal{H} = \frac{G_F \alpha V_{tb} V_{ts}^*}{\sqrt{2}\pi}\left[H_{1\mu}L^{\mu} + H_{2\mu}L^{5\mu}\right],$$

$$H_{1\mu} = C_9(\mu)\bar{s}\gamma_\mu(\mu)P_L b \; - \frac{2m_b}{q^2}C_7(\mu)\bar{s}i\sigma_{\mu\nu}q^\nu P_R b\,,$$

$$H_{2\mu} = C_{10}\bar{s}\gamma_\mu P_L b\,,\ L^\mu = \bar{l}\gamma^\mu l\,,\ L^{5\mu} = \bar{l}\gamma^\mu\gamma_5 l \qquad (1)$$

where V_{tq} $(q = s, b)$ are the CKM matrix elements, C_i $(i = 7, 9, 10)$ are the Wilson coefficients (WCs) and their expressions can be found in Ref. [11] for the SM. Since the operator associated with C_{10} is not renormalized under the QCD, it is the only one with the μ scale free. Besides the short-distance (SD) contributions, the main effect on the BR comes from $c\bar{c}$ resonant states such as Ψ, Ψ' *etc.*, *i.e.*, the long-distance (LD) contributions. In the literature [12], it has been suggested to combine the factorization assumption (FA) and vector meson dominance (VMD) approximation in estimating LD effects for the B decays. Hence, we may include the resonant effect (RE) by absorbing it to the related WC. The effective WC of C_9 is given by

$$C_9^{eff} = C_9(\mu) + (3C_1(\mu) + C_2(\mu))\Bigg(h(x,s) + \frac{3}{\alpha^2}\sum_{j=\Psi,\Psi'} k_j \frac{\pi\Gamma(j \to l^+l^-)M_j}{q^2 - M_j^2 + iM_j\Gamma_j}\Bigg)\,, \qquad (2)$$

where $h(x,s)$ describes the one-loop matrix elements of operators $O_1 = \bar{s}_\alpha\gamma^\mu P_L b_\beta\ \bar{c}_\beta\gamma_\mu P_L c_\alpha$ and $O_2 = \bar{s}\gamma^\mu P_L b\ \bar{c}\gamma_\mu P_L c$ [11], M_j (Γ_j) are the masses (widths) of intermediate states, and the factors $k_j \sim -1/(3C_1(\mu) + C_2(\mu))$ are phenomenological parameters for compensating the approximations of FA and VMD and reproducing the correct BRs $Br\,(B \to J/\Psi X \to l^+l^-X) =$

$Br\,(B \to J/\Psi X) \times Br\,(J/\Psi \to l^+l^-)$. Here, we have neglected the small Wilson coefficients. The transition amplitude for $B \to K^* l^+ l^-$ decays is found to be

$$\mathcal{M}^{(\lambda)}_{K^*} = \frac{G_F \alpha_{em} V_{tb} V^*_{ts}}{2\sqrt{2}\pi} \left\{ \mathcal{M}^{(\lambda)}_{1\mu} L^\mu + \mathcal{M}^{(\lambda)}_{2\mu} L^{5\mu} \right\}$$
$$\mathcal{M}^{(\lambda)}_{a\mu} = i f_1 \varepsilon_{\mu\nu\alpha\beta} \epsilon^{*\nu}(\lambda) P^\alpha q^\beta + f_2 \epsilon^*_\mu(\lambda) + f_3 \epsilon^* \cdot q P_\mu \quad (3)$$

where subscript $a = 1(2)$ while $f_i = h_i\ (g_i)\ (i = 1,\ 2,\ 3)$ and

$$h_1 = C_9^{eff}(\mu) V(q^2) - \frac{2m_b}{q^2} C_7(\mu) T(q^2),$$
$$h_{2(3)} = -C_9^{eff}(\mu) A_{0(1)}(q^2) + \frac{2m_b}{q^2} C_7(\mu) T_0(q^2),$$
$$g_1 = C_{10} V(q^2)\,, \quad g_{2(3)} = -C_{10} A_{0(1)}(q^2)\,. \quad (4)$$

Here, $P = p_1 + p_2$, $q = p_1 - p_2$, and the definitions of the form factors in Eq. (4) and the correspondences between our notations and those used in the literature can be found in the Appendix of Ref. [13].

To obtain the T-odd terms of $\varepsilon_{\mu\nu\alpha\beta} q^\mu \epsilon^{*\nu}(\lambda) p_l^\alpha P^\beta$, we have to study the processes of $B \to K^* l^+ l^- \to (K\pi) l^+ l^-$ so that the polarization λ and λ' in the differential decay rate, written as $d\Gamma \propto H(\lambda, \lambda')\ \mathcal{M}^{(\lambda)}_{K^*}\ \mathcal{M}^{(\lambda')\dagger}_{K^*}$ with $H(\lambda, \lambda') \equiv \epsilon(\lambda) \cdot p_K\ \epsilon^*(\lambda') \cdot p_K$, can be different. From Eq. (3), we see that $\mathcal{M}^{(\lambda)}_{2\mu}$ only depends on C_{10}. Clearly, the T violating effects can not be generated from $\mathcal{M}^{(\lambda)}_{2\mu} \mathcal{M}^{(\lambda')\dagger}_{2\mu'}$, but induced from $\mathcal{M}^{(\lambda)}_{1\mu} \mathcal{M}^{(\lambda')\dagger}_{1\mu'}$ and $\mathcal{M}^{(\lambda)}_{1\mu} \mathcal{M}^{(\lambda')\dagger}_{2\mu'}$. This can be understood as follows: firstly, for the $\mathcal{M}^{(\lambda)}_{1\mu} \mathcal{M}^{(\lambda')\dagger}_{1\mu'} Tr L^\mu L^{\mu'}$ contributions with $Tr L^\mu L^{\mu'} \sim \left(p^\mu_{l-} p^{\mu'}_{l+} + p^{\mu'}_{l-} p^\mu_{l+} - g^{\mu\mu'} p_{l-} \cdot p_{l+} \right)$, the relevant T-odd terms can be roughly expressed by

$$\begin{aligned}
&\mathcal{M}^{(\lambda)}_{1\mu} \mathcal{M}^{(\lambda')\dagger}_{1\mu'} Tr L^\mu L^{\mu'} \propto \\
&Z_1 Im h_1 h_3^* \epsilon(0) \cdot q \varepsilon_{\mu\nu\alpha\beta} q^\mu \epsilon^{*\nu}(\pm) p^\alpha_{l+} P^\beta + \\
&Z_2 Im h_1 h_2^* \epsilon(0) \cdot p_{l+} \varepsilon_{\mu\nu\alpha\beta} q^\mu \epsilon^{*\nu}(\pm) p^\alpha_{l+} P^\beta + \\
&Z_3 Im h_1 h_2^* \epsilon(\mp) \cdot p_{l+} \varepsilon_{\mu\nu\alpha\beta} q^\mu \epsilon^{*\nu}(\pm) p^\alpha_{l+} P^\beta
\end{aligned} \quad (5)$$

where $Z_i\ (i = 1, 2, 3)$ are constants. From Eq. (4), one gets $Im h_1 h_2^* \sim Im h_1 h_3^* \sim Im C_9^{eff}(\mu) C_7(\mu)$. We note that as shown in Eq. (5), the T-odd observables could be non-zero if the process involves strong phases or absorptive parts even without CP violating phases. By means of Eq. (2), $C_9^{eff}(\mu)$ includes the absorptive parts such that the results of Eq. (5) are not vanished in the SM. Secondly, for $\mathcal{M}^{(\lambda)}_{1\mu} \mathcal{M}^{(\lambda')\dagger}_{2\mu'} Tr L^\mu L^{5\mu'}$, one gets

$$\begin{aligned}
&\left(\mathcal{M}^{(\lambda)}_{1\mu} \mathcal{M}^{(\lambda')\dagger}_{2\mu'} + \mathcal{M}^{(\lambda)}_{2\mu} \mathcal{M}^{(\lambda')\dagger}_{1\mu'} \right) Tr L^\mu L^{5\mu'} \propto \\
&\left(Im h_2 g_3^* - Im h_3 g_2^* \right) \varepsilon_{\mu\nu\alpha\beta} q^\mu \epsilon^{*\nu}(\pm) p^\alpha_{l+} P^\beta
\end{aligned} \quad (6)$$

where $Tr L^\mu L^{5\mu'} = -4i\varepsilon^{\mu\mu'\alpha\beta} q_\alpha p_{l+\beta}$ has been used. From Eq. (4), we find that $Im h_2 g_3^* - Im h_3 g_2^*$ is only related to $Im C_7(\mu) C_{10}^*$ and the dependence of $Im C_9(\mu) C_{10}^*$ is canceled in Eq. (6). For the decays of $b \to s l^+ l^-$, since usually there are no absorptive parts in $C_7(\mu)$ and C_{10}, a non-vanishing value of $Im C_7(\mu) C_{10}^*$ indicates the pure weak CP violating effects.

In order to derive the whole differential decay rate for the K^* polarization, we choose $\epsilon(0) = (|\vec{p}_{K^*}|, 0, 0, E_{K^*})/m_{K^*}$, $\epsilon(\pm) = (0, 1, \pm i, 0)/\sqrt{2}$, and $p_{l^+} = \sqrt{q^2}(1, \sin\theta_l, 0, \cos\theta_l)/2$ with $E_{K^*} = (m_B^2 - m_{K^*}^2 - q^2)/2\sqrt{q^2}$ and $|\vec{p}_{K^*}| = \sqrt{E^2_{K^*} - m^2_{K^*}}$ in the q^2 rest frame and $p_K = (1, \sin\theta_K \cos\phi, \sin\theta_K \sin\phi, \cos\theta_K) m_{K^*}/2$ in the K^* rest frame where ϕ denotes the relative angle of decaying plane between $K\pi$ and l^+l^-. The differential decay rate with the only relevant terms is given by

$$\begin{aligned}
&\frac{d\Gamma}{d\cos\theta_K d\cos\theta_l d\phi dq^2} = \frac{3\alpha^2_{em} G_F^2 \left|\lambda_t\right|^2 |\vec{p}|}{2^{14}\pi^6 m_B^2} Br(K^* \to K\pi) \\
&\times \{ 4\cos^2\theta_K \sin^2\theta_l \sum_{i=1,2} |\mathcal{M}^0_i|^2 + \sin^2\theta_K (1 + \cos^2\theta_l) \\
&\sum_{i=1,2} \left(|\mathcal{M}^+_i|^2 + |\mathcal{M}^-_i|^2 \right) - \sin 2\theta_K \sin 2\theta_l \sin\phi \\
&\sum_{i=1,2} Im \left(\mathcal{M}^+_i - \mathcal{M}^-_i \right) \mathcal{M}^{0*}_i - 2\sin^2\theta_K \sin^2\theta_l \sin 2\phi \\
&\sum_{i=1,2} Im \left(\mathcal{M}^+_i \mathcal{M}^{-*}_i \right) + 2\sin 2\theta_K \sin\theta_l \sin\phi (Im \mathcal{M}^0_1 \\
&(\mathcal{M}^{+*}_2 + \mathcal{M}^{-*}_2) - Im(\mathcal{M}^+_1 + \mathcal{M}^-_1) \mathcal{M}^{0*}_2) + \cdots] \}\,,
\end{aligned} \quad (7)$$

where $|\vec{p}| = [((m_B^2 + m_{K^*}^2 - q^2)/(2m_b))^2 - m^2_{K^*}]^{1/2}$ and $\mathcal{M}^0_i$ and $\mathcal{M}^\pm_i$ denote the longitudinal and transverse polarizations of K^* respectively and their explicit expressions are as follows:

$$\mathcal{M}^0_a = \sqrt{q^2} \left(\frac{E_{K^*}}{m_{K^*}} f_2 + 2\sqrt{q^2} \frac{|\vec{p}_{K^*}|^2}{m_{K^*}} f_3 \right),$$
$$\mathcal{M}^\pm_a = \sqrt{q^2} \left(\pm 2 \left|\vec{p}_{K^*}\right| \sqrt{q^2} f_1 + f_2 \right),$$

The detailed derivation will be discussed elsewhere [14]. Other distributions for the K^* polarization can be found in Ref. [15]. From Eqs. (5) and (6), we know that $Im(\mathcal{M}^+_i - \mathcal{M}^-_i)\mathcal{M}^{0*}_i$ and $Im(\mathcal{M}^+_i \mathcal{M}^{-*}_i)$ are from $\mathcal{M}^{(\lambda)}_{1\mu} \mathcal{M}^{(\lambda')\dagger}_{1\mu'} Tr L^\mu L^{\mu'}$ while $Im \mathcal{M}^0_1 (\mathcal{M}^{+*}_2 + \mathcal{M}^{-*}_2) - Im(\mathcal{M}^+_1 + \mathcal{M}^-_1)\mathcal{M}^{0*}_2$ is induced by $\mathcal{M}^{(\lambda)}_{1\mu} \mathcal{M}^{(\lambda')\dagger}_{2\mu'} Tr L^\mu L^{5\mu'}$.

As seen from Eqs. (5) and (6), there are two possible sources for T violation, which are related to $Im C_9^{eff} C_7^*$ and $Im C_7 C_{10}^*$, respectively. In this paper, we only concentrate on the contribution from $Im C_7 C_{10}^*$ and explore the possibility of existing new CP violating phases. To do this, we examine the T-odd observable, defined by

$$\langle \mathcal{O} \rangle = \int \mathcal{O} d\Gamma\,. \quad (8)$$

where $\mathcal{O}$ is a T-odd five-momentum correlation, given by

$$\mathcal{O} = \frac{(\vec{p}_B \cdot \vec{p}_K)\,(\vec{p}_B \cdot (\vec{p}_K \times \vec{p}_{l^+}))}{|\vec{p}_B|^2\,|\vec{p}_K|^2\,\omega_{l^+}} \tag{9}$$

with $\omega_{l^+} = q \cdot p_{l^+}/\sqrt{q^2}$. In the K^* rest frame, we note that $\mathcal{O} = \cos\theta_K \sin\theta_K \sin\theta_l \sin\phi$. The statistical significance of the observable in Eq. (8) can be determined by

$$\varepsilon(q^2) = \frac{\int \mathcal{O} d\Gamma}{\sqrt{(\int d\Gamma)(\int \mathcal{O}^2 d\Gamma)}}\,. \tag{10}$$

Integrating all relevant angles in Eq. (10), we get

$$\begin{aligned}
\varepsilon(q^2) &\simeq \frac{0.76}{\sqrt{\mathcal{D}_1\mathcal{D}_2}}[Im\mathcal{M}_1^0(\mathcal{M}_2^{+*} + \mathcal{M}_2^{-*}) - \\
&\quad Im(\mathcal{M}_1^+ + \mathcal{M}_1^-)\mathcal{M}_2^{0*}]\,, \\
\mathcal{D}_1 &= \sum_{i=1,2}\left[\left|\mathcal{M}_i^0\right|^2 + \left|\mathcal{M}_i^+\right|^2 + \left|\mathcal{M}_i^-\right|^2\right]\,, \\
\mathcal{D}_2 &= \sum_{i=1,2}\left[\left|\mathcal{M}_i^0\right|^2 + \frac{1}{2}\left(\left|\mathcal{M}_i^+\right|^2 + \left|\mathcal{M}_i^-\right|^2\right)\right]\,.
\end{aligned} \tag{11}$$

To observe the effect at the $n\sigma$ level, the required number of B mesons is $N_B = n^2/(Br \cdot \varepsilon^2)$.

It is known that to study the exclusive decays of $B \to K^* l^+ l^-$, the main uncertainty is from the transition form factors in Eq. (4). The calculations of matrix elements for exclusive hadron decays can be performed in the perturbative QCD (PQCD) approach developed by Lepage-Brodsky (LB) [16]. However, with the LB approach, it has been pointed out that perturbative evaluation of the pion form factor suffers singularities from the end-point region with a momentum fraction $x \to 0$ [17]. In order to take care of the end-point singularities, the strategy of introducing k_T, the transverse momentum of the valence quark, and threshold resummations has been proposed and shown that the end-point singularities can be dealt with self-consistent in the PQCD [19]. Our calculations will base on such an approach. We use the results that have been displayed in Ref. [13]. The form factors given by the other QCD approaches such as the quark model (QM) and light-cone QCD sum rule (LCSR) can be found in Ref. [3,20]

It is well known that supersymmetric theories not only supply an elegant mechanism for the breaking of the electroweak symmetry and a solution to the hierarchy problem, but provide many new weak CP violating phases. Unfortunately, it has been shown that with the universal soft breaking parameters, these phases are severely bounded by electric dipole moments (EDMs) of electron and neutron [21] so that the contributions to ϵ and ϵ' become negligible. However, one may avoid the EDM constraints by setting the squark masses of the first two generations to be as heavy as few TeV [22]. The SUSY models with the non-universal soft A terms inspired by string theories [23] and left-right symmetry [24], which have unsuppressed weak CP phases and lead to the observed ϵ, ϵ' and the large CPA of $B \to X_s\gamma$, have also been proposed.



FIG. 1. The differential decay BRs as a function of s: The solid (dashed), long dash-dotted (dotted) and dash-dotted (dense-dotted) lines describe the results of the PQCD (LCSR) in the standard, model-independent with $ImC_7 = 0.25$ and SUSY models, respectively.

To illustrate the new physics contributions, we shall use (a) a model-independent approach by assuming that the CPV only arises from C_7, the main effect for $b \to s\gamma$, with taking $ImC_7 = 0.25$ and the values of remaining WCs are the same as those in the SM; and (b) the results of the generic supersymmetric extension of the SM in Ref. [25] where, instead of scanning the whole allowed parameter space, we take the values

$$\begin{array}{ll}
Re\,C_7^{SUSY} \simeq 0, & Im\,C_7^{SUSY} \simeq -0.27, \\
Re\,C_9^{SUSY} \simeq -0.6, & ImC_9^{SUSY} \simeq -0.1, \\
Re\,C_{10}^{SUSY} \simeq 8.5, & Im\,C_{10}^{SUSY} \simeq -\,2.5\,.
\end{array} \tag{12}$$

Here, the scale for WCs without specified is M_W, which can be taken approximately the same as the SUSY scale. Several remarks concerning on the SUSY models are given as follows: (i) the source of the CP violating phases is embedded in the sfermion mass matrices with the mass-insertion method [25]; (ii) we assume that the flavor diagonal terms in the SUSY models are either real or small so that there are no new constraints from the EDMs of neutron and electron; (iii) for $Im\ C_7^{SUSY} = \pm 0.27$, we find that $Br(B \to X_s\gamma)$ increases $\sim$ 10% by comparing with that in the SM [26], consistent with the data of $(3.22 \pm 0.40) \times 10^{-4}$ [27], and the rate CP asymmetry in $b \to s\gamma$ is about $\pm 4\%$, much larger than the SM prediction of $\sim$ 1% [28] but still within the recent 95% range of -0.30 to 0.14 implied by the CLEO measurement [29]; and (iv) the SUSY effect on $B \to J/\Psi\,K$ with the values in Eq. (12) is small, while those with other allowed parameters in SUSY have been discussed in Ref. [7].

With the values in Eq. (12), we find that the BRs of $B \to K^* l^+ l^-$ in the standard, model-independent and SUSY models are 1.33 (1.88), 1.51 (2.06) and 1.29 (5.17) $\times 10^{-6}$ for the PQCD (LCSR), respectively. We note that the corresponding BRs for $B \to K l^+ l^-$ are 5.32 (5.17), 5.36 (5.22) and 5.19 (4.94) $\times 10^{-7}$, respectively, which so far are all consistent with the experimental data [2]. The differential decay rates of $B \to K^* l^+ l^-$



FIG. 2. The statistical significance ε of the T violating observable $\langle\mathcal{O}\rangle$ in (a) the model-independent approach with $ImC_7 = 0.25$ and (b) the generic SUSY model as a function of s. The solid and dash-dotted lines stand for the results of the PQCD and LCSR, respectively.

as a function of $s = q^2/m_B^2$ are shown in Fig. 1, and the distribution of $\varepsilon(q^2)$ is displayed in Fig. 2. From Fig. 1, we see that the uncertainty from the different QCD approaches is compatible to the new physics effects. Thus, it is not easy to ensure the existence of new physics by only measuring the branching distribution. However, the T violating observable $\langle\mathcal{O}\rangle$ which is zero in the SM, could be large. In particular, in the SUSY model, the statistical significance $\varepsilon(q^2)$ of $\langle\mathcal{O}\rangle$ can be over 30% at the low q^2 region and it is insensitive to non-PQCD effects.

Finally, we note that the origin of the T violating effect here would be quite different from the CPA in $b \to s\gamma$. For example, $\varepsilon(q^2)$ can be non-zero and large even for the case of $Im\,C_7 = 0$ with CPV arising from phases in non-dipole WCs such as C_{10} due to new physics, whereas $a_{CP}(b \to s\gamma)$ is the same as that in the SM.

In summary, we have shown that, by measuring the angular distributions of K and leptons, one can obtain the individual informations of longitudinal and transverse polarizations of K^* which are all sensitive to physics beyond the SM. For reducing the uncertainties of QCD effects, one can define some physical observables normalized by the differential decay rate. Among different angular distributions, we have found that the T-odd contribution arising from $M_{1\mu}^{(\lambda)} M_{2\mu'}^{(\lambda')\dagger} Tr L^{\mu} L^{5\mu'}$ is purely related to the weak CP violating phase, which could be sizable in new physics such as the model with SUSY. Searching for such a T violating distribution, one can distinguish the new CP violating source from the CKM mechanism. We remark that to observe the T violating effects of $\langle\mathcal{O}\rangle$ in $B \to K^* l^+ l^-$ $(l = e, \mu)$ with $\varepsilon \sim 10\%$ at the 1σ level, at least 5×10^7 B mesons are needed, which can be done at the B-factories in KEK and SLAC and future hadron colliders.

Acknowledgments: This work was supported in part by the National Science Council of the Republic of China under Contract Nos. NSC-89-2112-M-007-054 and NSC-89-2112-M-006-033.

---

[1] CLEO Collaboration, M. S. Alam et. al., Phys. Rev. Lett. **74**, 2885 (1995).
[2] Belle Collaboration, K. Abe et. al., hep-ex/0109026.
[3] For a recent review, see A. Ali et. al., *Phys. Rev.* **D61**, 074024 (2000).
[4] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
[5] Belle Collaboration, H. Yamamoto, talk presented at the fifth KEK topical conference, Nov. 20-22, 2001.
[6] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. 87, 091801 (2001); BELLE Collaboration, A. Abashian *et al.*, Phys. Rev. Lett. 87, 091802 (2001).
[7] For a recent review, see E. Lunghi and D. Wyler, hep-ph/0109149; and references therein.
[8] R. Garisto, Phys. Rev. D**51**, 1107 (1995); C.H. Chen, C.Q. Geng and C.C. Lih, *ibid*, D**56** 6856 (1997).
[9] C.Q. Geng and C.P. Kao, Phys. Rev. D**57**, 4479 (1998); C.H. Chen and C.Q. Geng, *ibid*, D**64** 074001 (2001).
[10] F. Krüger and E. Lunghi, Phys. Rev. D**63** 014013 (2000).
[11] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys **68,** 1230 (1996).
[12] N.G. Deshpande, J. Trampetic and K. Panose, Phys. Rev. D**39**, 1462 (1989); C.S. Lim, T. Morozumi, and A.T. Sanda, Phys. Lett. B**218**, 343 (1989); A. Ali, T. Mannel, and T. Morozumi, *ibid*, B**273**, 505 (1991); F. Krüger and L.M. Sehgal, *ibid*, B**380**, 199 (1996).
[13] C.H. Chen and C.Q. Geng, Phys. Rev. D**63**, 114025 (2001).
[14] C.H. Chen and C.Q. Geng, hep-ph/0203003.
[15] C.S. Kim, *et.al.* Phys. Rev. D**62**, 034013 (2000); T. M. Aliev, *et.al.*, Phys. Lett. B**511**, 49 (2001).
[16] P. Lepage and S. Brodsky, Phys. Lett. B**87**, 959 (1979); Phys. Rev. D**22**, 2157 (1980).
[17] N. Isgur and C.H. Llewellyn-Smith, Nucl. Phys. B**317**, 526 (1989); A. Szczepaniak, E. M. Henley, and S. Brodsky, Phys. Lett. B**243**, 287 (1990); R. Akhoury, G. Sterman and Y.P. Yao, Phys. Rev. D**50**, 358 (1994).
[18] M. Beneke and T. Feldmann, Nucl. Phys. B**592**, 3 (2000).
[19] H-n. Li, Phys. Rev. D**64**, 014019 (2001).
[20] T. M. Aliev, *et.al.* Phys. Rev. D**56**, 4260 (1997); D. Melikhov and B. Stech, *ibid*, D**62**, 014006 (2000).
[21] R. Garisto and J.D. Wells, Phys. Rev. D**55**, 1611 (1997); R. Garisto, *ibid*, D**49** 4820 (1994) and the references therein.
[22] K.S. Babu, *et.al.*, Phys. Rev. D**59**, 016004 (1999).
[23] D. Bailin and S. Khalil, Phys. Rev. Lett. **86**, 4227 (2001)
[24] K.S. Babu, *et.al.*, Phys. Rev. D**61**, 091701 (2000).
[25] F. Gabbiani, *et.al.* Nucl. Phys. B**477**, 321 (1996); E. Lunghi, *et.al. ibid.* **568,** 120 (2000).
[26] A. L. Kagan and M. Neubert, Eur. Phys. J. C**7**, 5 (1999).
[27] S. Ahmed *et al.* [CLEO Collaboration], hep-ex/9908022; R. Barate *et al.* [ALEPH Collaboration], Phys. Lett. B **429**, 169 (1998); K. Abe *et al.* [Belle Collaboration], *ibid*, B **511**, 151 (2001).
[28] A. L. Kagan and M. Neubert, Phys. Rev. D**58**, 094012 (1998).
[29] T. E. Coan *et al.* [CLEO Collaboration], Phys. Rev. Lett. **86**, 5661 (2001).